

REVEALING THE NATURE OF THE HIGHLY OBSCURED GALACTIC SOURCE IGR J16318–4848⁰

S. Chaty,^{1,2,4} and P. Filliatre^{2,3}

RESUMEN

El resumen será traducido al español por los editores. The X-ray source IGR J16318–4848 was the first source discovered by INTEGRAL on 2003, January 29. We carried out optical and near-infrared (NIR) observations at the European Southern Observatory (ESO La Silla) in the course of a Target of Opportunity (ToO) programme. We discovered the optical counterpart and confirmed an already proposed NIR candidate. NIR spectroscopy revealed a large amount of emission lines, including forbidden iron lines and P-Cygni profiles. The spectral energy distribution of the source points towards a high luminosity and a high temperature, with an absorption greater than the interstellar absorption, but two orders of magnitude lower than the X-ray absorption. We show that the source is an High Mass X-ray binary (HMXB) at a distance between ~ 1 and ~ 6 kpc, the mass donor being an early-type star, probably a sgB[e] star, surrounded by a rich and absorbing circumstellar material. This would make the second High Mass X-ray Binary (HMXB) with a sgB[e] star after CI Cam, indicating that a new class of strongly absorbed X-ray binaries is being unveiled by INTEGRAL.

ABSTRACT

The X-ray source IGR J16318–4848 was the first source discovered by INTEGRAL on 2003, January 29. We carried out optical and near-infrared (NIR) observations at the European Southern Observatory (ESO La Silla) in the course of a Target of Opportunity (ToO) programme. We discovered the optical counterpart and confirmed an already proposed NIR candidate. NIR spectroscopy revealed a large amount of emission lines, including forbidden iron lines and P-Cygni profiles. The spectral energy distribution of the source points towards a high luminosity and a high temperature, with an absorption greater than the interstellar absorption, but two orders of magnitude lower than the X-ray absorption. We show that the source is an High Mass X-ray binary (HMXB) at a distance between ~ 1 and ~ 6 kpc, the mass donor being an early-type star, probably a sgB[e] star, surrounded by a rich and absorbing circumstellar material. This would make the second High Mass X-ray Binary (HMXB) with a sgB[e] star after CI Cam, indicating that a new class of strongly absorbed X-ray binaries is being unveiled by INTEGRAL.

Key Words: STARS: CIRCUMSTELLAR MATTER — STARS: EMISSION-LINE — BE—X-RAYS: BINARIES — IGR J16318-4848

IGR J16318–4848 has been the first source to be discovered by the INTEGRAL imager IBIS/ISGRI on 2003 January 29, 0.5° south from the galactic equator (Courvoisier et al. 2003). A subsequent observation by XMM-Newton on 2003 February 10 localized it with an accuracy of $4''$, the high energy spectrum suggesting a high column density of $N_H > 10^{24} \text{ cm}^{-2}$ (Matt & Guainazzi (2003), Walter et al. (2003)). In the course of a ToO programme at ESO to look for counterparts of high energy sources discovered by satellites including IN-

TEGRAL (PI S. Chaty), we carried out on 2003, February, 23-25 photometric and spectroscopic observations in the optical and NIR of the high-energy source IGR J16318–4848, with EMMI and SOFI instruments on ESO/NTT. We discovered the optical counterpart and confirmed the NIR one (see Walter et al. (2003)) by independent astrometry. The optical/NIR images and spectra are shown in Figures 1, 2, 3, 4 and 5. We derived the absorption along the line of sight: $A_V \sim 17.4$ magnitudes, and the temperature of the companion star: ~ 18000 K. The distance of the source is constrained between ~ 1 and ~ 6 kpc. The $0.95 - 2.52 \mu\text{m}$ NIR spectrum is highly unusual, very rich in emission lines, suggesting a highly complex and stratified circumstellar environment, or an envelope. Study of the spectral lines suggest a sgB[e] star so the system would be

¹Université Paris 7, 2 place Jussieu, F-75 005 Paris, France.

²Service d'Astrophysique, DSM/DAPNIA/SAP, CEA/Saclay, F-91 191 Gif-sur-Yvette, Cedex, France.

³Fédération de Recherche APC, Paris, France.

⁴chaty@cea.fr

⁰Based on observations collected at the European Southern Observatory, Chile (observing proposal ESO N° 70.D-0340).




Fig. 1. R band image of the field of view of IGR J16318-4848. We reported the XMM uncertainty circle of 4".




Fig. 2. Ks band image of the same field.

a HMXB, probably hosting a neutron star, like CI Cam. The reader should consult Filliatre & Chaty (2004) for more details. INTEGRAL is on the course of revealing a new population of obscured high energy sources, which might help us to understand the evolution of high-energy binary systems.

SC is grateful to Rob Hynes, Joanna Mikolajewska, Ignacio Negueruela and Marc Ribó for very helpful discussions on the nature of the source during this IAU colloquium.

REFERENCES

- Courvoisier, T.J.-L., Walter, R., Rodriguez, J. et al. 2003, IAU Circ., 8063
 Filliatre, P., Chaty, S., 2004, ApJ subm.
 Matt, G. & Guainazzi, M. 2003, MNRAS, 341, L13
 Walter, R., Rodriguez, J., Foschini, L. et al. 2003, A&A, 411, L427




Fig. 3. NIR spectrum (0.95–1.65 μm)




Fig. 4. NIR spectrum (1.5–2.05 μm)




Fig. 5. NIR spectrum (2.0–2.55 μm)